

Recent Results from the Pegasus Toroidal Experiment¹

A.J. REDD, J.L. BARR, M.W. BONGARD, M.G. BURKE, R.J. FONCK, E.T. HINSON, D.J. SCHLOSSBERG, N.L. SCHOENBECK, K.E. THOME, Univ of Wisc - Madison — Pegasus is an ultra-low aspect ratio spherical tokamak ($A=1.15$), used to study the physics of low- A plasmas and develop non-solenoidal tokamak startup techniques. A combination of point-source magnetic helicity injection and poloidal field induction produces non-solenoidal tokamak plasmas with $I_p \leq 0.17$ MA using 4 kA injected current I_{inj} , consistent with the helicity injection rate and a Taylor relaxation limit. A strong double layer sheath at the injectors describes the impedance of the injection circuit, implying the helicity injection rate for a given I_{inj} is a strong function of the local plasma density. Impurity ion spectroscopy indicates strong heating ($T_i \approx 0.5$ keV) during helicity drive. Passive gas-fueled electrodes can be used as helicity injectors, complementing active current sources and affording a means to optimize the Taylor and helicity injection limits. Ohmic plasmas in Pegasus are often unstable to peeling modes, an instability underlying edge localized modes (ELMs) in larger tokamaks. Time-resolved current profile measurements show that ELM-like, current-carrying filaments form at the plasma edge, which then detach and propagate outward.

¹Work supported by US DOE OFES, grant DE-FG02-96ER54375

- Prefer Oral Session
 Prefer Poster Session

Aaron Redd
aredd@wisc.edu
University of Wisconsin - Madison