

Low-m Tearing Mode Studies on Pegasus¹ J.L. BARR, M.W. BONGARD, R.J. FONCK, B.T. LEWICKI, A.J. REDD, University of Wisconsin-Madison — Large scale, low-m/n=1 tearing activity is commonly observed in ohmic discharges on PEGASUS. Accentuated by broad regions of low magnetic shear, this tearing activity can limit performance in high TF utilization regimes ($I_p/I_{TF} \approx 1$) where q_0 is a low-order rational. In addition to increasing TF to raise q_0 , tearing activity can be mitigated by: TF ramp-downs; electrostatic preionization to obtain OH breakdown at lower TF; and, DC helicity injection to provide strong edge current drive. Systematic scans of TF, I_p , and $\partial I_p/\partial t$ are presently being conducted to characterize these modes under a range of operational conditions. Measurements of magnetic activity via internal Mirnov probes are complicated by electrostatic noise arising from PEGASUS' high-frequency switching power supplies. This noise can be mitigated somewhat by shielding and signal processing, but can be eliminated altogether by freewheeling all power supplies for a brief period.

¹Work supported by US DOE Grant DE-FG02-96ER54375.

- Prefer Oral Session
 Prefer Poster Session

Michael Bongard
mbongard@wisc.edu
University of Wisconsin-Madison

Special instructions: Please place as poster 6/7 with the other Pegasus presentations. JLB's membership is pending, and starts 8/1/10.