

Effect of Aspect Ratio on H-mode and ELM Characteristics¹ K.E. THOME, G.M. BODNER, M.W. BONGARD, M.G. BURKE, R.J. FONCK, D.M. KRIETE, University of Wisconsin-Madison — The H-mode confinement regime is achieved at near-unity aspect ratio ($A < 1.2$) in the Pegasus Toroidal Experiment via high-field-side fueling and low edge recycling. Ohmic H-mode is attained in both limited and diverted magnetic topologies. This regime is characterized by: reduced D_α emissions; increased core rotation; increased central heating; formation of edge current and pressure pedestals; and measured energy confinement consistent with the ITER98pb(y,2) scaling. The H-mode power threshold, P_{LH} , behaves quite differently at low- A when compared with high- A operations. P_{LH}/P_{LH_ITPA08} increases sharply as A is lowered and no difference in P_{LH} for limited and diverted plasmas is observed at $A \sim 1.2$. No minimum in P_{LH} with density is observed. Some of these results are consistent with the FM³ model for the L-H transition.² Two classes of ELMs have been observed. Small, Type III-like ELMs are present at low input power and have $n \leq 4$. At $P_{OH} \gg P_{LH}$, they transition to large, Type-I-like ELMs with intermediate $5 < n < 15$. The Type III ELM magnetic structures behave opposite that of high- A plasmas, with n much higher, presumably due to the naturally higher J/B peeling mode drive at low- A . Long-sought measurements of the $J_{edge}(R, t)$ pedestal collapse during an ELM event show a complex, multimodal pedestal collapse and the subsequent ejection of a current-carrying filament.

¹Work supported by US DOE grant DE-FG02-96ER54375.

²Fundamenski *et al.*, Nucl. Fusion **52**, 062003 (2012.)

Prefer Oral Session

Prefer Poster Session

K.E. Thome

kethome@wisc.edu

University of Wisconsin-Madison

Special instructions: Please place as poster 3 within the Pegasus presentations.